



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

12th of June 173 deaths have been registered. Of these 1 was from scarlet fever, 5 from measles, 4 from diphtheria, 1 from smallpox, and 5 from typhoid fever.

SPIRIDION C. ZAVITZIANO,
United States Sanitary Commissioner.

STATISTICAL REPORTS.

AUSTRALIA—*New South Wales—Sydney.*—Month of April, 1898. Estimated population, 417,250. Total number of deaths, 434, including diphtheria, 12; enteric fever, 10; measles, 3, and scarlet fever, 3.

BERMUDA.—Two weeks ended June 17, 1898. Estimated population, 15,013. Total number of deaths, 1.

CHILE—*Antofagasta.*—Month of May, 1898. Estimated population, 14,000. Total number of deaths, 46, including 2 from phthisis pulmonalis.

FRANCE—*Rouen.*—Month of May, 1898. Estimated population, 112,657. Total number of deaths, 250, including enteric fever, 1; measles, 1; whooping cough, 1, and 53 from phthisis pulmonalis.

GREAT BRITAIN—*England and Wales.*—The deaths registered in 33 great towns in England and Wales during the week ended June 18 correspond to an annual rate of 14.7 a thousand of the aggregate population, which is estimated at 11,218,378. The highest rate was recorded in New Castle on Tyne, viz, 22.2, and the lowest in Cardiff, viz, 8.5.

London.—One thousand one hundred and ninety-three deaths were registered during the week, including measles, 50; scarlet fever, 10; diphtheria, 26; whooping cough, 44; enteric fever, 4, and diarrhea and dysentery, 12. The deaths from all causes correspond to an annual rate of 13.8 a thousand. In greater London 1,550 deaths were registered, corresponding to an annual rate of 12.6 a thousand of the population. In the "outer ring" the deaths included 11 from measles, 10 from diphtheria, 3 from scarlet fever, and 11 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended June 18 in the 23 principal town districts of Ireland was 19.6 a thousand of the population. The lowest rate was recorded in Portadown, viz, 0.0, and the highest in Newtownards, viz, 45.4 a thousand. In Dublin and suburbs 147 deaths were registered, including enteric fever, 4, and whooping cough, 1.

Scotland.—The deaths registered in 8 principal towns during the week ended June 18 correspond to an annual rate of 20.2 a thousand of the population, which is estimated at 1,568,536. The lowest mortality was recorded in Paisley, viz, 17.2, and the highest in Perth, viz, 27.2 a thousand. The aggregate number of deaths registered from all causes was 609, including measles, 32; scarlet fever, 8, and whooping cough, 22.